

Table of contents *(in brief)*

| | |
|--|------------|
| Chapter 1: Biosynthesis, production, quality and stability of pigments | 1 |
| Chapter 2: Food processing | 81 |
| Chapter 3: Pigments from biotechnology (microorganisms, microalgae, plant cell cultures)..... | 175 |
| Chapter 4: Biological activity, bioavailability, and toxicology..... | 219 |
| Chapter 5: Composition, characterization and analytical aspects of pigments and their derivatives | 299 |
| Late posters (titles only) | 414 |

Note from the Editor:

Papers submitted to publication into the Proceedings were peer-reviewed by the members of the PF2004 Scientific committee.

“Peer review is a referring process used to check the quality and importance of reports of research. An article submitted for publication in a peer-reviewed book is reviewed by other experts in the area. It aims to provide a wider check on the quality and interpretation of a short report and to improve its quality” (*The Cochrane Library Glossary, 1996*).

Table of contents

□ Chapter One

Biosynthesis, production, quality and stability of pigments

| | |
|---|----|
| Regulation of carotenoid biosynthesis in orange fruits (<i>Citrus sinensis</i> L. Osbeck) Rodrigo M.J., Alqu zar B., Marcos J.F., Zacar as L. | 2 |
| <i>In Planta</i> active anthocyanin degradation in Brunfelsia flowers Vaknin H., Ovadia R., Nissin-Levi A., Oren-Shamir M. | 5 |
| Chlorophylls and their breakdown products in green plant tissues: Genetic and environmental control Roca M., James C., H rtensteiner S., Thomas H., Ougham H. | 8 |
| Identification and functional characterisation of rat liver bilitranslocase in carnation petals Passamonti S., Cocolo A., Braidot E., Petrusa E., Peresson C., Medic N., Macr  F., Vianello A. | 11 |
| Coloured Swiss chard (<i>Beta vulgaris</i> L. ssp. <i>cicla</i> [L.] Alef. cv. "Bright Lights") - a new source of betalains ? Stintzing F.C., Kugler F., Carle R. | 14 |
| Fluorescent pigments. New perspectives in betalain research and applications Gand a-Herrero F., Garc a-Carmona F., Escribano J. | 17 |
| More than just colours: Carotenoid pigments are key precursors for aroma compounds in tomato fruit Lewinsohn E., Azulay Y., Bar E., Meir A., Hirschberg J., Zamir D., Tadmor Y. | 20 |
| Carotenoids and enzymatic aroma formation in nectarines Baldermann S., Naim M., Winterhalter P., Fleischmann P. | 23 |
| Recovery of grape pomace anthocyanins. Influence of the maceration time during winemaking G mez-Plaza E., Mi ano A., L pez-Roca J.M. | 26 |
| Utilisation of supercritical fluid techniques in the fractionation of natural pigments J rvenp   E.P., Rostiala N., Huopalahti R. | 29 |
| Influence of enzyme maceration on anthocyanin extraction from blackcurrant <i>Ribes nigrum</i> (L.) berries Lapornik B., Pro ek M., Petri  D., Golc-Wondra A. | 32 |
| Kinetic and thermodynamic measurements of flavylum hydration of the major diacylated anthocyanin isolated from sweet potato (<i>Ipomoea batatas</i> [L.] Lam.) Malien-Aubert C., Pauleau N., Ben Hassam K., Dangles O., Jouenne E. | 35 |

| | |
|---|----|
| Application of tristimulus colorimetry to optimize the extraction of anthocyanins from jaboticaba (<i>Myrciaria jaboticaba</i> Berg.) fruits | |
| Montes J.C., Raymundo M., Vicario I.M., Fett R., Heredia F.J. | 38 |
| Viable alternative processings to increase xanthophyll extraction from marigold flower (<i>Tagetes erecta</i>) | |
| Navarrete-Bolaños J.L., Jimenez-Islas H., Botello-Alvarez E., Rico-Martinez R. | 41 |
| Impact of various factors on the composition and stability of blackcurrant anthocyanins | |
| Rubinskiene M., Viskelis P., Jasutiene I., Bobinas C. | 44 |
| Yellow highly polar pigment prepared from apple dihydrochalcones | |
| Sanoner P., Guyot S., Sanogo T., Lequeré J-M., Drilleau J-F., Renard C.M.G.C. | 47 |
| Co-polymerization and co-pigmentation: effects on anthocyanin stability | |
| Wesche-Ebeling P., Argaíz-Jamet A., Olmedo-Hernández M.M., Mayoral G. | 50 |
| Light stability of spray-dried encapsulated bixin with different edible polysaccharide preparations | |
| Barbosa M. I. J., Borsarelli C. D., Mercadante A. Z. | 53 |
| Stability of black carrot anthocyanins in various juices | |
| Kirca A., Özkan M., Cemeroğlu B. | 56 |
| Relationships between chromatic characteristics of red wines and their different fractions of anthocyanin compounds | |
| Pérez-Magariño S., González-SanJosé M.L. | 59 |
| Comparison of methods for the deodorization of turmeric | |
| Silva L. V., Glória M.B.A. | 62 |
| Xanthophylls of vegetables – the most common carotenoids in nature : identification, localization and conformation <i>in vivo</i> | |
| Ruban A.V. | 65 |
| New syntheses of natural carotenoid isorenieratene (ϕ,ϕ-carotene) | |
| Valla A., Cartier D., Valla B., Le Guillou R., Andriamialisoa Z., Breithaupt D., Labia R., Dufossé L. | 67 |
| Photosensitized degradation of all-<i>trans</i>-lycopene | |
| Nunes I.L., Borsarelli C.D., Mercadante A.Z. | 70 |
| Bistable aggregate of all-<i>trans</i>-β-carotene in an aqueous solution | |
| Mori Y. | 72 |
| Phenolic colourants obtained by enzymatic synthesis with a fungal laccase in hydro-organic biphasic system | |
| Mustafa R., Muniglia L., Rovel B., Girardin M. | 75 |
| Adsorption and recovery of anthocyanins on resins | |
| Scordino M., Di Mauro A., Passerini A., Maccarone E. | 78 |

□ Chapter Two

Food processing

| | |
|---|-----|
| Characterization of commercial olive oils from their pigment visible light absorption bands at 610–720 nm following treatment with bleaching earths Ayuso J., Ruiz A., Escolar D., Haro M ^a R..... | 82 |
| Storage influence on the initial content and class of pigments of virgin olive oil Gallardo-Guerrero L., Roca M., Gandul-Rojas B., Mínguez-Mosquera M.I. | 85 |
| Individual behavior of olive fruits during table olive processing Gandul-Rojas B., Jarén-Galán M., Mínguez-Mosquera M.I. | 88 |
| Conventional and non conventional olive processing techniques affect lipochromes and other virgin oil biomarkers Ranalli A., Malfatti A., Lucera L., Contento S., Pavone G. | 91 |
| Changes in colour and pigments in spinach after cooking and drying Kidmose U., Christensen L.P., Edelenbos M. | 94 |
| “Maillard cross-links” formed within soy protein isolate films slows down fat oxidation of high omega 3 fish oil Beng G.T., Miskin M.N., Bee C.P., Easa A.M..... | 97 |
| Colouring potential differences between melanoidins formed in absence and in presence of phenolic compounds Ortega M., González-Sanjosé M ^a L., Muñiz P. | 100 |
| Quantitative and structural changes of betacyanins in red beet (<i>Beta vulgaris</i> L.) juice induced by thermal treatment Herbach K.M., Stintzing F.C., Carle R..... | 103 |
| Effects of dehydration on the antioxidant activities and anthocyanin contents of Saskatoon berries (<i>Amelanchier alnifolia</i> Nutt) Kitts D.D., Hu C., Kwok B., Durance T..... | 106 |
| Degradation of various fruit juice anthocyanins by hydrogen peroxide Özkan M., Yemenicioğlu A., Cemeroglu B. | 109 |
| Extraction and use of anthocyanins from maize (<i>Zea mays</i>) grains as colouring in yoghurt Salinas-Moreno Y., Díaz Velázquez A., Rubio-Hernández D. | 112 |
| Effect of pH, sugar and heating temperature on the anthocyanin patterns and antioxidant capacity of mulberry extract Tsai P.-J., Yu T.-Y., Huang Y.-T. | 115 |
| Phycocyanin and Lutein coloured food emulsions – relation between pigment concentration and structural properties Batista A.P., Raymundo A., Sousa I., Empis J..... | 118 |
| Colouring emulsions using microalgal biomass - stability studies Gouveia L., Raymundo A., Batista A.P., Miranda A., Sousa I., Empis J..... | 121 |
| Thermal colour degradation kinetics of date paste Ahmed J..... | 124 |

| | |
|--|-----|
| The colour of <i>Brevibacterium linens</i> and <i>Arthrobacter</i> species depends on the deacidification yeast and the cheese-making technology Leclercq-Perlat M.N., Corrieu G., Spinnler H.E. | 127 |
| Effect of the physical state on carotenoid oxidation in tomato derivatives Manzocco L., Calligaris S., Nicoli M.C. | 130 |
| Lycopene from tomato product added minced meat. Effect on storage quality and colour Østerlie M., Lerfall J. | 133 |
| Effect of pasteurization on the carotenoid composition of Valencia orange juice Gama J.J.T., Sylos C.M. | 136 |
| Impact of thermal processing on retinol equivalent values of paprika oleoresins Pérez-Gálvez A., Jarén-Galán M., Mínguez-Mosquera M. I. | 139 |
| <i>Trans-cis</i>-isomerization of β-carotene in carrot juices and functional drinks Schieber A., Marx M., Carle R. | 142 |
| Muscle of smoked rainbow trout (<i>Onchorhynchus mykiss</i>) packaged under modified atmosphere. Evolution of colour, pH, exudate, oxidative process and microbiological aspect Quiñones A., Choubert G., Gómez R. | 145 |
| Effect of light on colour stability of sliced smoked rainbow trout <i>Oncorhynchus mykiss</i> fed astaxanthin Choubert G., Dentella E., Atgié C., Baccaunaud M. | 147 |
| Content of α- and β-carotene in raw and fried carrots of different colours Kidmose U., Christensen L.P., Edelenbos M., Bjørn G. K. | 151 |
| Spray drying microencapsulation of the bixin natural colourant Constant P. B. L., Stringheta P. C., Coimbra J. S. R., Sandi D. | 154 |
| Effect of the concentration process on carotenoid composition of Valencia orange juice Gama J.J.T., Sylos C.M. | 157 |
| Effect of thermal dehydration of spice red pepper on the stability of carotenoids and antioxidants Daood H.G., Kapitány J., Albrecht K., Biacs P. | 160 |
| Comparison of three methods for aroma generation from apocarotenoids and oleoresin isolated from Euphorbiaceae (physical, enzymatic and microbiological processes) Del Toro C. L., Waché Y., Villanueva S., Lugo E. | 163 |
| <i>Euterpe oleracea</i> juice as functional pigment for yogurt Coisson J.D., Travaglia F., Piana G., Capasso M., Arlorio M. | 166 |
| A porcine pigment (AproRed[®]) to improve colour in cooked hams Oliver M.A., Polo J., Panella N., Arnau J., Gil M., Contreras M., Morera S., Ruíz J. | 169 |

| | |
|--|-----|
| Thermal damage on pigments of blood orange juice and advantages of reverse osmosis application | |
| Di Mauro A., Arena E., Fallico B., Scordino M., Maccarone E..... | 172 |
| | |
| □ Chapter Three | |
| <i>Pigments from biotechnology</i> | |
| <i>(microorganisms, microalgae, plant cell cultures...)</i> | |
| | |
| Plant cell cultures and micro algae for the production of natural colourants for food application | |
| Ravishankar G.A. | 176 |
| | |
| Simple extraction of carotenoids from red yeast, <i>Xanthophyllomyces dendrorhous</i> | |
| Song J.-Y., Kwak W.-K., An G.-H..... | 179 |
| | |
| Carotenoid extraction with supercritical CO₂ from a marine strain of <i>Synechococcus</i> sp. | |
| Lama C.M., Macias-Sánchez M.D., Lubián L.M., Mantell C., Rodríguez M., Martínez de la Ossa E., Montero O. | 182 |
| | |
| Studies on <i>Xanthophyllomyces dendrorhous</i> cultivated in batch fermentation | |
| Martin A.M., Sivagurunathan M. | 186 |
| | |
| Factors that control gamete differentiation in the astaxanthin producing microalgae <i>Haematococcus pluvialis</i> | |
| Martín L., Vargas M.A., Cid A., Magnúsdóttir R., Norsker N.-H., Rodríguez H..... | 189 |
| | |
| Methods for astaxanthin extraction from <i>Haematococcus pluvialis</i> | |
| Zou N., Li B., Yi J., Zhang X., Sun D..... | 192 |
| | |
| Unique food colours from algae: fluorescent pink and marine blue | |
| Yaron A., Arad S. (Malis) | 195 |
| | |
| <i>Monascus</i> yellow pigments, more than colours... | |
| Yongsmith B..... | 196 |
| | |
| Astaxanthin production by an extremely halophilic archaea isolated from high salt fermented Thai foods, <i>Halobacterium</i> sp. BCC12460 | |
| May T. B., Lumyong S., Sukchotiratana M., Paukatong K..... | 198 |
| | |
| Microalgal β-carotene in diets enhances astaxanthin levels in shrimp <i>Litopenaeus vannamei</i> | |
| Paniagua-Michel J., Yepiz-Velazques R. | 201 |
| | |
| <i>Chlorella vulgaris</i> biomass as fat mimetic in oil-in-water food emulsions | |
| Raymundo A., Gouveia L., Batista A.P., Resendes M., Empis J., Sousa I. | 204 |
| | |
| Arpink Red – meet a new natural red food colourant of microbial origin | |
| Sardaryan E., Zihlova H., Strnad R., Cermakova Z. | 207 |
| | |
| Production of β-carotene by a <i>Sphingomonas</i> mutant | |
| Silva C., Cabral J.M.S., van Keulen F..... | 209 |
| | |
| Partially saturated xanthophylls from <i>Aspergillus carbonarius</i> | |
| Vijayalakshmi G., Venkatesh K.S., Vanajakshi V., Kumaresan N., Ravi-Kumar K., Umesh-Kumar S. | 212 |

| | |
|--|-----|
| Antimicrobial activities of <i>Monascus</i> pigments derived with various amino acids Kim C., Jung H., Shin C.S..... | 215 |
|--|-----|

□ **Chapter Four**

Biological activity, bioavailability and toxicology

| | |
|--|-----|
| Antioxidant and biological activity of phenolic pigments from <i>Theobroma cacao</i> hulls extracted with supercritical CO₂ Arlorio M., Coïsson J.D., Travaglia F., Varsaldi F., Miglio G., Lombardi G., Martelli A. .. | 220 |
| Antioxidant properties of model Maillard systems prepared from glucose with proline, glycine, lysine, arginine, cysteine or glutathione Billaud C., Peyrat-Maillard M-N., Chow Y-N., Berset C., Nicolas J. | 223 |
| Effect of Maillard reaction conditions on the increase in free radical scavenging activity of browning products obtained from tuna waste hydrolysate Sumaya-Martinez M.T., Guerard F., Thomas S., Linard B., Binet A. | 226 |
| Colour and antioxidant activity of melanoidins of some Spanish Coffee Liqueurs Greco L., Fogliano V., González-Sanjósé M ^a L..... | 230 |
| Antioxidant activities of berry anthocyanins Kähkönen M., Viljanen K., Heinonen M. | 233 |
| Antioxidant activity of chlorophylls and their derivatives by the β-carotene bleaching method Lanfer-Marquez U.M., Barros R.M.C., Sinnecker P..... | 235 |
| Study on sea buckthorn (<i>Hippophae rhamnoides</i> L.) pigments as potential antioxidants Lõugas T., Vokk R. | 238 |
| Evaluation of the TEAC method for antioxidant capacity determination of azorubine Obón J.M., Castellar M.R., Cascales J.A., Fernández-López J.A. | 241 |
| Antioxidant molecules and free radical processes in carrot hypercotyl root stock – variations with position, age and post-harvest storage time Pirker K.F., Goodman B.A., Nagreiter C., Reichenauer T.G..... | 244 |
| Optimization of the antioxidant capacity of a mixture of carotenoids and α-tocopherol in the development of a nutritional supplement Yoshimura K., Castro I.A., Barros S.B.M., Marques U. L., Motizuki M., Sawada T. C. H. | 246 |
| Turmeric, a natural pigment : From food preservation to health promotion Nagabhushan M., Kolpe U., Ramaswamy V., Rao B.S.S. | 249 |
| Antigenotoxic and antimutagenic potential of an annato pigment (norbixin) against oxidative stress Júnior A. C. T. S., Asad L. M. B. O., de Oliveira E. B., de Souza G. P., Kovary K., Asad N. R., Felzenszwalb I. | 252 |

| | |
|---|-----|
| Anthocyanins and bioactivities of black rice (<i>Oryza sativa</i> L. <i>Indica</i>) Hu C., Zawistowski J., Kitts D.D. | 255 |
| Survey of usage and estimated intakes of annatto extracts Tennant D. R, O’Callaghan M. | 258 |
| An <i>in vitro</i> model to study the intestinal absorption of carotenoids During A., Harrison E. H. | 261 |
| Intestinal absorption of lycopene and isomer pattern in human plasma as affected by the food matrix Kaufmann K., Fröhlich K., Bitsch R., Böhm V. | 265 |
| Thermal processing of carrots : lycopene stability and isomerisation in regard to bioavailability Mayer-Miebach E., Behsnilian D., Regier M., Schuchmann H.P. | 268 |
| A pharmacokinetic evaluation of the urinary excretion of dietary betalains following consumption of red beet juice Netzel M., Frank T., Straß G., Quaas D., Stintzing F.C., Carle R., Bitsch R., Bitsch I. | 271 |
| Tissue distribution of anthocyanins following gastric absorption of a pure anthocyanin mixture Passamonti S., Vrhovsek U., Vanzo A., Mattivi F. | 274 |
| Postprandial evolution of the carotenoid content in the TRL fraction after a single ingestion of virgin olive oil in humans Pérez-Gálvez A., Pacheco Y.M., Bermúdez B., López S., Abia R., Muriana F.J.G., Villar J., Garrido-Fernández J. | 277 |
| Uptake of lutein by human intestinal Caco-2 TC-7 cells partly occurs through a facilitated process involving the scavenger receptor class B type I (SR-BI) Reboul E., Abou L., Mikail C., Ghiringhelli O., André M., Portugal H., Jourdeuil-Rahmani D., Lairon D., Amiot-Carlin M.-J., Borel P. | 280 |
| Sunlight rapidly raises lycopene and β-carotene at the site of erythema in human skin Richelle M., Offord E.A., Liardet S., Hager C., Bortlik K.H., Tavazzi I., Saucy F. , Duruz E., Hohlfeld P., Halliwell B., Applegate L. | 283 |
| Dose-dependent urinary excretion of dietary anthocyanins following ingestion of elderberry juice Straß G., Netzel M., Frank T., Dietrich H., Bitsch I., Bitsch R. | 286 |
| Absorption and distribution in erythrocytes and low density lipoproteins of betalains from cactus pear (<i>Opuntia ficus indica</i>) in healthy humans. Potential health effects of the dietary betalains Tesoriere L., Butera D., Allegra M., Livrea M.A. | 288 |
| Effect of tomato product consumption on the plasma status of antioxidant microconstituents and on the plasma total antioxidant capacity in healthy subjects Tyssandier V., Feillet-Coudray C., Caris-Veyrat C., Guillard J.C., Coudray C., Bureau S. , Reich M., Amiot-Carlin M.J., Bouteloup-Demange C., Boirie Y., Borel P. | 291 |
| Bioavailability of lycopene isomers from sauces prepared from different tomato varieties Nuray Unlu Z., Clinton S. K., Francis D. M., Schwartz S. J. | 295 |

□ Chapter Five

Composition, characterization and analytical aspects of pigments and their derivatives

| | |
|--|-----|
| Colour characteristics of Monastrell wines depending on the date of harvest Bautista-Ortín A.B., López-Roca J.M., Fernández-Fernández J.I., Gómez-Plaza E..... | 300 |
| The effect of 3-O-β-glucosylation on the pH-dependent absorption spectra and structural transformations of anthocyanins Borkowski T., Szymusiak H., Gliszczyńska-Swigło A., Tyrakowska B..... | 303 |
| Oxidative cleavage products of lycopene: production, characterization and mechanism of formation Caris-Veyrat C., Schmid A., Carail M., Böhm V..... | 306 |
| Differential partitioning of betalains in aqueous two phase extraction Chethana S., Nayak C. A., Raghavarao K.S.M.S..... | 309 |
| Characterization and application of betalain pigments from plants in the Amaranthaceae Corke H., Cai Y.Z., Sun M..... | 312 |
| High-performance liquid chromatography of carotenoids from tomato products on conventional and base-deactivated AB reversed phase column Daood H.G., Biacs P., Samer H. | 315 |
| Carotenoid composition of two Brazilian acerola genotypes (<i>Malpighia puniceifolia</i> L.) De Rosso V. V., Mercadante A. Z..... | 318 |
| Varietal and interspecific influence on carotenoid content in citrus from the Mediterranean area Dhuique-Mayer C., Caris-Veyrat C., Ollitrault P., Curk F., Amiot-Carlin M.J..... | 321 |
| Spectrocolorimetry in the CIE L*a*b* colour space as useful tool for monitoring the ripening process and the quality of red-smear soft cheeses Dufossé L., Galaup P., Carlet E., Flamin C., Valla A..... | 323 |
| Anthocyanin composition of pigmented potato varieties Eichhorn S., Winterhalter P..... | 326 |
| Identification of novel anthocyanin copigmentation products in berry juices Eiro M.J., Heinonen M..... | 329 |
| The fingerprint of pigments on the fluorescence of virgin olive oil Escolar D., Haro M ^a R., Ayuso J..... | 331 |
| HPLC analysis of the pigments produced by the microfora isolated from ‘Protected Designation of Origin’ French red-smear soft cheeses Galaup P., Flamin C., Carlet E., Dufossé L..... | 334 |
| HPLC quantification of carotenoids in Valencia orange juice Gama J.J.T., Sylos C.M..... | 338 |

| | |
|---|-----|
| Oxygenated lycopene and dehydrated lutein in tomato puree Etoh H., Yokota T., Suzuki R., Yoshioka R., Inakakuma T., Ishiguro Y. | 341 |
| Routine and sensitive SPE-HPLC method for quantitative determination of pheophytin <i>a</i> and pyropheophytin <i>a</i> in olive oils Hornero-Méndez D., Gandul-Rojas B., Mínguez-Mosquera M.I..... | 344 |
| The use of reflectance spectroscopy to estimate carotenoid concentration in cereal grains Humphries J.M., Mares D.J., Graham R.D. | 347 |
| Intrinsic variability of olive fruits harvested for table olive processing Jarén-Galán M., Gallardo-Guerrero L., Mínguez-Mosquera M.I..... | 350 |
| Fast and simple analysis method for the anthocyanins isolated from various plant sources Järvenpää E.P., Parbo I., Huopalahti R. | 353 |
| Composition and properties of red grape marc pigments Jones G.P., Elieff S., Swinny E.E., Obradovic D., Roe S.J..... | 356 |
| Production of β-ionone from β-carotene through enzyme-generated radicals degradation in a biphasic medium Ly M.H., Waché Y., Belin J.-M. | 359 |
| Carotenoid content and colour of Valencia late ultrafrozen juices from southern Spain Meléndez-Martínez A.J., Britton G., Vicario I.M., Heredia F.J..... | 361 |
| Carotenoids profile during Port wine aging Mendes-Pinto M.M., Silva Ferreira A.C., Guedes de Pinho P. | 364 |
| Carotenoid components in yellow, green, and black soybean varieties and immature young soybean (Eda-mame) Momma M., Saito M., Chikuni K. | 367 |
| Astaxanthin in feeds and fishes : A global analytical approach Morice G., Le Texier S., Baudry F., Metra P.L..... | 370 |
| Analysis of a complex mixture of carotenes from oil palm (<i>Elaeis guineensis</i>) fruits Mortensen A. | 372 |
| Colour studies of model solutions containing betaxanthins and betacyanins Møbhammer M.R., Stintzing F.C., Carle R..... | 375 |
| Aggregation of natural food dye molecules and its spectroscopic display Nizamov N., Astanov S., Nizamov Sh. N. | 378 |
| Pigments in Colombian fruits : Anthocyanins from Corozo (<i>Bactris major</i>) fruit Osorio C., Hillebrand S., Duque C., Winterhalter P..... | 381 |
| Separation and characterization of curcuminoid pigments from <i>Curcuma longa</i> Péret-Almeida L., Alves J.R., Glória M.B.A..... | 384 |
| Carotenoids in different hybrids of Sea buckthorn berries Pintea A., Varga A., Socaciu C. | 387 |

| | |
|---|-----|
| Isolation of flavanol-anthocyanin adducts by countercurrent chromatography Salas E., Poncet-Legrand C., Fulcrand H., Meudec E., Köhler N., Winterhalter P. , Cheynier V. | 390 |
| Characterization of anthocyanins extracts from maize (<i>Zea mays</i>, L.) grains Rubio-Hernández D., Salas-Sánchez G., Salinas-Moreno Y., Ramos-Lobato N. A. | 393 |
| Anthocyanins and carotenoids in varieties of fruits and vegetables Sass-Kiss A., Kiss J., Milotay P., Kerek M., Toth M. | 396 |
| Comparative fruit colouration in watermelon and tomato Tadmor Y., King S., Levi A., Davis A., Hirschberg J. | 400 |
| Differential Scanning Calorimetry (DSC) to evaluate technological antioxidant activity of phenolic pigments from <i>Corylus avellana</i> perisperm Travaglia F., Capasso M., Coïsson J. D., Piana G., Arlorio M. | 403 |
| Tracking colour and pigment changes in anthocyanin products Wrolstad R. E. | 406 |
| Anthocyanins, chlorophylls and xanthophylls in pistachio nuts (<i>Pistacia vera</i>) of different geographic origin Bellomo M. G., Fallico B. | 408 |
| Carotenoid composition of camu-camu (<i>Myrciaria dubia</i>) from two Brazilian cities Zanatta C. F., Mercadante A. Z. | 411 |
| ❑ Late posters | 414 |